

Detecting Glandular lesions
by
Cervical Cytology

Dr Margaret Sage
March 2018

Glandular abnormalities reported in NZ in 2016

Cytology:

Atypical Glandular Cells/AIS = 403 reports (0.10%)

Adenocarcinoma (all types) = 101 reports (0.02%).

HSIL = 4347 reports (1.1%)

SCC = 35 reports (0.01%)

Histology:

Adenocarcinoma in situ (AIS) = 149 women

Endocervical adenocarcinoma = 19 women

HSIL = 4945 women

SCC = 140 women

The NZ Cervical Cancer Audit 2000 - 2002

- 336 smears from 178 women taken within 4 years prior to a histological diagnosis of invasive cervical cancer were rescreened.
- 18% of 160 negative smears prior to invasive SCC were upgraded to “high-grade”
- 22% of 65 negative smears prior to invasive (endocervical) adenocarcinoma were upgraded to “high-grade”

Overview



Endocervical cells

1. Normal/reactive endocervical cells
2. Atypical endocervical cells
3. Adenocarcinoma in situ (AIS)
4. Invasive endocervical adenocarcinoma



Endometrial cells

1. Normal endometrial cells
2. Atypical endometrial cells
3. Endometrial adenocarcinoma


Other abnormal glandular cells



Normal endocervical cells



Reactive endocervical cells



ThinPrep

Atypical glandular cells: AIS on follow-up


Adenocarcinoma in situ (AIS)

Cell aggregates: **ARCHITECTURE** matters most


- large irregularly shaped sheets
- tightly crowded cells with nuclear overlapping
- sheet edges: palisaded nuclei, common border, feathering
- strips and rosettes with pseudostratification
- gland openings
- papillary groups

Cell morphology:


- few single cells, but can see more in LBC samples
- hyperchromatic; mild increase in nuclear size
- chromatin uniform but stippled or granular
- mitoses, may be abnormal
- apoptotic debris




Adenocarcinoma in situ (AIS): Histology




AIS Histology: Endocervical type



AIS Histology: Endometrioid type



AIS Histology: Intestinal type




AIS: Irregular crowded groups




ThinPrep

AIS: Cell crowding and nuclear overlapping




SurePath

AIS: Common sheet edge




ThinPrep

AIS: Strip of a sheet edge




AIS: Rosette, mitoses, feathering




SurePath

AIS: Abnormal Strips




AIS: Abnormal strip




SurePath

AIS: More abnormal strips




ThinPrep

AIS: possible rosette



SurePath



AIS: Cell morphology




ThinPrep

AIS: Hyperchromasia


AIS: well-differentiated



AIS: poorly-differentiated



Endometrioid AIS



ThinPrep


Reported as Atypical Glandular Cells
Follow-up was AIS: intestinal type

Endocervical adenocarcinoma

1. Features of AIS


2. Features suggestive of invasion

- tumour diathesis
- single cells
- marked pleomorphism
- nuclei: chromatin clearing, conspicuous nucleoli
- fewer strips and rosettes
- supercrowding with loss of polarity




SurePath

Endocervical adenocarcinoma




ThinPrep


Endocervical adenocarcinoma




SurePath



ThinPrep




SurePath




ThinPrep


Differential Diagnoses: High sampling





Cervical Endometriosis




Tubal metaplasia



HSIL involving glands




Normal endometrial cells




Atypical endometrial cells (LBC)
Follow-up: Endometrial hyperplasia


Atypical endometrial cells



Endometrial Adenocarcinoma




Endometrial hyperplasia




Atypical endometrial cells

F/U: Well-differentiated endometrial adenocarcinoma



SurePath

Endometrial Adenocarcinoma




Endometrial Adenocarcinoma



ThinPrep

Endometrial Adenocarcinoma



Metastatic Gastric adenocarcinoma (LBC)